The A/C study: A cross-sectional study of HIV epidemiology among African, Caribbean and Black people in Ontario

Mbuagbaw L*, Tharao WE, Husbands W, Nelson LE, Aden M, Arnold K, Baidoobonso S, Dabone C, Dryden O, Etowa EB Hamid J, Jackson-Best F, Kohoun B, Lawson DO, Lofters A, Luyombya H, Mbulaheni T, Mkandawire P, Ndungu M, Nyambi A, Obiorah S, Ongoiba F, Ongolo-Zogo C, Oraka C, Shahin R, Yaya S, Hendriks A, Gebremeskel A, Inoua H, Etowa J

Conflict of Interest Disclosure:

*I have no conflicts of interest



African Caribbean Black Health Toronto • Ottawa



CAHR 2020 Virtual
May 1 & 2, 2020
Hope, Victories and Perseverance beyond 2020

I'ACRV 2020 virtuel sur votre agenda
1 au 2 mai, 2020
Espoir, Victoires et Persévérance au-delà de 2020



Background

African, Caribbean, and Black (ACB) communities are disproportionately infected by HIV in Ontario, Canada. They constitute only 5% of the population of Ontario yet account for 25% of new diagnoses of HIV.¹

Aims


The aim of this study is to understand underlying factors that augment the HIV risk in ACB communities and to inform policy and practice in Ontario.

Methods

- Design: Cross-sectional study of first- and second-generation ACB adults aged 15-64 in Toronto and Ottawa.
- Dates: Nov 2018 Dec 2019
- Following principles of community-based and community-led research.
- Strong stakeholder engagement
- Integrated knowledge translation
- Analysis: Descriptive statistics and Logistic regression



Community Based Research







Results

Table 1: Characteristics of participants

Variable	Statistic (N=1380)	
City: n (%)		
Toronto (or greater Toronto area)	854 (61.9)	
Ottawa (or greater Capital region)	526 (38.1)	
Age (years): n (%)	,	
15 – 19	157 (11.4)	
20 – 29	426 (30.9)	
30 – 39	338 (24.5)	
40 – 49	286 (20.7)	
50 – 59	108 (7.8)	
60 – 64	30 (2.2.)	
Gender identity: n (%)		
Man	481 (35.6)	
Woman	842 (61.8)	
Trans persons	6 (0.4)	
Non-binary persons	16 (1.1)	
Tested for HIV (Yes): n (%)	950 (68.8)	
HIV diagnosis (Self report)*	75 (8.1); 95% CI 6.4-10.1	
HIV diagnosis (Dried Blood Spot)*	67(8.4); 95% CI 6.7-10.3	
Born in Canada (Yes): n (%)	298 (21.6)	
Employment: n (%)		
Unemployed/Students	641 (46.4)	
Part time	262 (19.0)	
Full time	477 (34.6)	
Education: n (%)		
University	739 (55.7)	
College	249 (18.8)	
High school	309 (23.3)	
Less than high school	30 (2.3)	

*N=923; **N=942; CI: Confidence Intervals

Table 2: Factors associated with diagnosis HIV infection[¥]

Variables	OR (95% CI)	Р
Age	2.19 (1.57-3.05)	<0.001
Ever had an STI test	0.8 (0.35-1.81)	0.596
Employment		
Full time employment	1	
Unemployed	2.43 (0.99-5.94)	0.052
Part time employment	3.65 (1.24-10.69)	0.018
Gender		
Man	1	
Woman	0.58 (0.28-1.22)	0.153
Trans/non-binary persons	6.12 (1.00-37.88)	0.049
Born in Canada (Yes)	0.19 (0.04- 0.92)	0.039
Age at first intercourse	0.97 (0.9-1.05)	0.460
Level of Education		
University	1	
Less than high school	11.45 (1.8-72.83)	0.010
High school	2.12 (0.83-5.39)	0.116
College	2.17 (0.91-5.14)	0.079

 $^{^{*}}$ Based on Dried Blood Spot; STI: Sexually Transmitted Infection; OR: Odds Ratio; CI: Confidence Intervals; Model fit: Nagelkerke R^{2} =0.23

Toronto • Ottawa

Conclusions

- Among the ACB participants in this study, older people, people without full-time employment, and people who have less than a high school education, trans and non-binary participants are more likely to be diagnosed with HIV.
- People born in Canada are <u>less likely</u> to be diagnosed with HIV.

Contacts

Lawrence Mbuagbaw: mbuagblc@mcmaster.ca
Winston Husbands: whusbands@ohtn.on.ca
Wangari Tharao: wangari@whiwh.com

Josephine Etowa: Josephine. Etowa@uottawa.ca

Acknowledgements















Shannon Ryan Hector Addison

